

Rate each question from 0 to 5 based on the descriptions at the top of the numbered columns. Write your answer in the first open column to the right. Total your answers to get your final score. Then take this symptom journal to your doctor to interpret the results.

<b>Over the past month, how often have you...</b>	<i>Not at all</i>	<i>Less than 1 time in 5</i>	<i>Less than half the time</i>	<i>About half the time</i>	<i>More than half the time</i>	<i>Almost always</i>	Date	Date	Date	Date	Date	Date
Had a sensation of not emptying your bladder completely after you finished urinating?	0	1	2	3	4	5						
Had to urinate again less than 2 hours after you finished urinating?	0	1	2	3	4	5						
Stopped and started again several times as you urinated?	0	1	2	3	4	5						
Found it difficult to postpone urination?	0	1	2	3	4	5						
Had a weak urinary stream?	0	1	2	3	4	5						
Had to push or strain to begin urination?	0	1	2	3	4	5						
Typically needed to get up to urinate from the time you went to bed at night until the time you got up in the morning?	0	1	2	3	4	5						
							Total	Total	Total	Total	Total	Total

UROXATRAL is an  $\alpha_1$ -blocker for the treatment of the signs and symptoms of BPH.

**Important Safety Information**

BPH and prostate cancer can cause the same symptoms. However, UROXATRAL is not a treatment for prostate cancer.

**Please see additional Important Safety Information on back and accompanying important product information.**



*Powerful relief. Simply.*

**Bother Score due to urinary symptoms  
(Do not add to total)**

	<i>Delighted</i>	<i>Pleased</i>	<i>Mostly satisfied</i>	<i>Mixed</i>	<i>Mostly dissatisfied</i>	<i>Unhappy</i>	<i>Terrible</i>	Date	Date	Date	Date	Date	Date
How would you feel if you had to live with your current urinary condition—no better, no worse—for the rest of your life?	0	1	2	3	4	5	6						

**Additional Important Safety Information**

Do not take UROXATRAL if you have liver problems or if you are taking antifungal drugs like ketoconazole or itraconazole, or HIV drugs like ritonavir.

UROXATRAL can cause a sudden drop in blood pressure, especially when starting treatment. This may lead to fainting, dizziness, and lightheadedness. Do not drive, operate machinery, or do any dangerous activity until you know how UROXATRAL will affect you. This is especially important if you already have a problem with low blood pressure or take medicines to treat high blood pressure. There may be an increased risk of low blood pressure and fainting when taking UROXATRAL in combination with blood pressure medication or nitrates, or erectile dysfunction medication.

If considering cataract surgery (clouding of the eyes), tell your eye surgeon that you are currently taking UROXATRAL or have previously been treated with an alpha-blocker.

Before taking UROXATRAL, tell your doctor if you have kidney problems.

Also, tell your doctor if you or any family member(s) have or take medications for a rare heart condition known as congenital prolongation of the QT interval.

The most common side effects with UROXATRAL are dizziness, upper respiratory tract infection, headache, and tiredness.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

**Please see accompanying important product information.**

